

TriElements® Clinical Reference Guide

**MacroMicro™ Cleanse plus
ReEssence Cellulite Support**

Awarded 4 patents**By the U.S. Government

Scientific Advanced Complete
Dietary Supplement 120 Capsules

MacroMicro Cleanse

Directions: Take 2 capsules daily for maintenance and 4 capsules daily periodically, when a more vigorous cleans/detox is needed. Best taken with food or a large glass of water in divided doses (am & pm). Store in a cool dry place

Serving Size: 2 Vegetarian Capsules (vegetable cellulose)
Servings Per Container 30

	Amount Per Serving	%DV
Milk Thistle Extract (seed) (80% silymarin)	150 mg	*
N-Acetyl Cysteine	125 mg	*
Broccoli extract (aerial) (4000 ppm Sulforaphane)	40 mg	*
Alpha Lipoic Acid	60 mg	*
Proprietary Blend: CranXact® Extract** (cranberry, grape, apple) (Standardized to clinical levels of bacteria anti-adhesion activity) blended with special detox/cleansing support ingredients.	545 mg	*
Dandelion (root)		*
Cranberry Extract (seeds and/or fruit)		*
Burdock (root)		*
Grape Extract (seeds)		*
Pomegranate Extract (fruit) (40% Ellagic Acid)		*
Artichoke Extract (leaf) (2% Cynarin)		*
Red Clover (aerial)		*
Mullein (leaf)		*
Ginger (root)		*
Larch Arabinogalactan		*
Apple Extract (skin, seeds, fruit)		*

*Daily Value not established

Other Ingredients: vegetable cellulose, silica

**CranXact U.S. patent #'s 5,474,774, 5,525,341, 5,646,178, 5,650,432. Bacteria anti-adhesion activity levels based on red blood cell hamagglutination assay.

ReEssence Cellulite Support

Directions: Take 2 capsules daily. When more support is needed take 2 capsules twice daily. Best taken with food or a glass of water. Store in a cool dry place

Serving Size: 2 Capsules
Servings per container 30

	Amount Per Serving	%DV
Vitamin C (ascorbic acid, plant source capsicum frutescens fruit)	120 mg	200%
Pantothenic Acid (as calcium pantothenate)	20 mg	200 %
Riboflavin (vitamin B2)	13 mg	765%
Niacin (Vitamin B3)	6 mg	30%
Iodine (plant source Fucus vesiculosus)	57 mcg	38 %
Centella Asiatica Extract (leaf) (standardized to 20% Asiaticosides)	100 mg	*
Bromelain (standardized to 2000 GDU)	250 mg	*
Lemon Bioflavonoids	100 mg	*
Milk Thistle Extract (seed) (standardized to 80% silymarin)	88 mg	*
Butcher's Broom Extract (root) (standardized to 10% Ruscogenins)	50 mg	*
Capsicum frutescens (fruit) (minimum 40,000 heat units)	40 mg	*
Green Tea Extract (leaf) (standardized to 95% catechins, 50% EGCG))	20 mg	*

*Daily Value (DV) not established

Purpose of Formula:

Cleanse & Detox

Comprehensive & Advanced: This combination product is uniquely patented and formulated to provide comprehensive support for detoxification and cleansing at the macrobiological and microbiological levels and provide a more balanced approach to the body's phase 1 and phase 2 detoxification systems. It can be used to support daily detox or a short term detox/cleanse program

Macrobiological: It is specially designed to help all of the systems and organs of the body in their detoxification and cleansing functions. It also helps strengthen these systems and organs so they are more capable of performing the detoxification/cleansing functions on their own.

Microbiological: It inhibits the adherence of bacteria to living tissue so they can be safely and naturally voided from the body. It works at the cellular level to enhance energy production in the mitochondria and inhibit associated oxidation. It protects cells from alteration and damage.

Cellulite Nutrient Support

Cellulite tissue has poor circulation and is starved for nutrients partly because it is laden with toxic, stagnant waste which also damages the tissue and blood vessels. In this condition nutrients are blocked from reaching cellulite tissue and can't provide the nourishment that is needed. Combining detoxification and waste removal with special cellulite specific nutrients helps provide a more effective and lasting improvement to health and appearance. The cellulite specific nutrients provided in this formulation powerfully combat cellulite laden tissue in these four areas:

- **Fat** – Supports healthy metabolism & the conversion of fat into energy

- **Connective Tissue** – Promotes re-growth of healthy, supple connective tissue.
- **Circulation** – Helps tone and strengthen blood vessels and maintain healthy microcirculation

Application and Key Benefits:

- Designed to provide a balanced approach to Phase I and Phase II detoxification support. (1,2,3,4)
- Supports the detoxification functions of the liver and also promotes the natural production and recycling of glutathione, a very important endogenous detoxifier and antioxidant. It also protects the liver by inhibiting passage of toxins into liver cells and stimulating regeneration of new liver cells. (5,6,7,8,10,11,12)
- Supports normal bile flow and fat metabolism which helps nutrient absorption and elimination of waste. (9)
- Helps to improve the efficiency of energy production in the mitochondria and prevent associated oxidative cellular damage.(19,20,21,22,23,27,)
- Inhibits Glycation and ‘Glycation End Products’ (AGE’s). (28,29, 30, 31, 32)
- Protects cells from alteration and damage. (10,11,12)
- Supports healthy respiration by helping to thin stubborn mucous (13)
- Helps the body detoxify and protect against chemicals and heavy metal toxins such as mercury and arsenic. (14,15)
- Anti-oxidant support for both fat and water soluble systems (16)
- Provides high levels of bacteria anti-adhesion support (17,18)
- Helps regenerate and recycle Vitamins C, E, Glutathione, and Coenzyme Q10 (20,21,22,23)
- Helps support and protect normal structure and function for blood, colon and kidneys. (24,25,26)
- Helps maintain normal connective tissue structure & function in the skin. (33,34,35,36)
- Helps burn fat for energy, tone and strengthen blood vessels and maintain a healthy microcirculation and fluid balance (37-53)

These statements have not been evaluated by the food and drug administration. This product is not intended to diagnose, cure, treat, or prevent disease

1. Valenzuela A, et al. Selectivity of Silymarin on the Increase of the Glutathione Content in Different Tissues of the Rat. *Planta Medica*. 1989;55:1550-52.
2. Beloqui O, et al. N-acetyl cysteine enhances the response to interferon-alpha in chronic hepatitis C: a pilot study. *J Interferon Res*. Aug1993;13(4):279-82.
3. Fahey JW, Talalay P. Antioxidant functions of sulforaphane: a potent inducer of Phase II detoxication enzymes. *Food Chem Toxicol* 1999;37:973-9.
4. Maheo K, Morel F, Langouet S, et al. Inhibition of cytochromes P-450 and induction of glutathione S- transferases by sulforaphane in primary human and rat hepatocytes. *Cancer Res* 1997;57:3649-52.
5. Flora K, et al. Milk Thistle (*Silybum marianum*) for the Therapy of Liver Disease. *Am J Gastroenterol*. 1998;93(2):139-43.
6. Salmi H, et al. Effect of Silymarin on Chemical, Functional, and Morphological Alterations of the Liver. A Double-blind Controlled Study. *Scand J Gastroent*. 1982;17:517-21.
7. Sonnenbichler J, Zetl I. Stimulating influence of a flavonolignan derivative on proliferation, RNA synthesis and protein synthesis in liver cells. In *Assessment and Management of Hepatobiliary Disease*, ed. L Okolicsanyi, G Csomos, G Crepaldi. Berlin: Springer-Verlag, 1987, 265-72.
8. Beloqui O, et al. N-acetyl cysteine enhances the response to interferon-alpha in chronic hepatitis C: a pilot study. *J Interferon Res*. Aug1993;13(4):279-82.
9. *Cardui mariae fructus* (Milk Thistle fruit). Commission E Monograph. Mar1986;Bundesanzeiger:no. 50.
- 10 Campos R, et al. Silybin Dihemisuccinate Protects Against Glutathione Depletion and Lipid Peroxidation Induced by Acetaminophen on Rat Liver. *Planta Medica*. 1989;55:417-19.
11. Rue YC. Advances in Pharmacological Studies of Silymarin. *Mem Inst Oswaldo Cruz*. 1991;86(Suppl 2):79-85.
12. Dehmlow C, Erhard J, de Groot H. Inhibition of Kupffer Cell Functions as an Explanation for the Hepatoprotective Properties of Silibinin. *Hepatology*. Apr1996;23(4):749-54.
13. Olsson B, et al. Pharmacokinetics and bioavailability of reduced and oxidized N-acetylcysteine. *Eur J Clin Pharmacol* 1988;34:77-82.
14. Flora SJ, et al. Arsenic-induced oxidative stress and its reversibility following combined administration of N-acetylcysteine and meso 2,3-dimercaptosuccinic acid in rats. *Clin Exp Pharmacol Physiol*. Nov1999;26(11):865-9.
15. Ballatori N, et al. N-acetylcysteine as an antidote in methylmercury poisoning. *Environ Health Perspect*. May1998;106(5):267-71.
16. Kagan VE, Shvedova A, Serbinova E, et al. Dihydrolipoic acid--a universal antioxidant both in the membrane and in the aqueous phase. Reduction of peroxy, ascorbyl and chromanoxyl radicals. *Biochem Pharmacol*. Oct1992;44(8):1637-49.
17. Avorn J, Monane M, Gurwitz JH, et al. Reduction of bacteriuria and pyuria after ingestion of cranberry juice. *JAMA* 1994;271:751-4.
18. Zafiri D, Ofek I, Adar R, et al. Inhibitory activity of cranberry juice on adherence of type 1 and type P fimbriated *Escherichia coli* to eucaryotic cells. *Antimicrob Agents Chemother* 1989;33:92-8.
19. Hagen TM, Ingersoll RT, Lykkesfeldt J, Liu J, Wehr CM, Vinarsky V, Bartholomew JC, Ames AB. (R)-alpha-lipoic acid-supplemented old rats have improved mitochondrial function, decreased oxidative damage, and increased metabolic rate. *FASEB J*. 1999 Feb;13(2):411-8.
20. Busse E, Zimmer G, Schopohl B, et al. Influence of alpha-lipoic acid on intracellular glutathione in vitro and in vivo. *Arzneimittel-Forschung*. 1992;42:829-831.
21. Packer L, Witt EH, Tritschler HJ. alpha-Lipoic acid as a biological antioxidant. *Free Radic Biol Med*. Aug1995;19(2):227-50.
22. Stoyanovsky DA, Goldman R, Darrow RM, et al. Endogenous ascorbate regenerates vitamin E in the retina directly and in combination with exogenous dihydrolipoic acid. *Curr Eye Res*. Mar1995;14(3):181-9.
23. Kagan V, Serbinova E, Packer L. Antioxidant effects of ubiquinones in microsomes and mitochondria are mediated by tocopherol recycling. *Biochem Biophys Res Comm*. 1990;169:851-857
24. Gavish D, et al. Lipoprotein (a) Reduction by N-Acetylcysteine. *Lancet*. Jan1991;337:203-204.
25. Shyu KG, Cheng JJ, Kuan P. Acetylcysteine protects against acute renal damage in patients with abnormal renal function undergoing a coronary procedure. *J Am Coll Cardiol* 2002;40:1383-8.
26. Estensen RD, Levy M, Klopp SJ, et al. N-acetylcysteine suppression of the proliferative index in the colon of patients with previous adenomatous colonic polyps. *Cancer Lett* 1999;147:109-14.
27. Hagen TM, Ingersoll RT, Lykkesfeldt J, Liu J, Wehr CM, Vinarsky V, Bartholomew JC, Ames AB. (R)-alpha-lipoic acid-supplemented old rats have improved mitochondrial function, decreased oxidative damage, and increased metabolic rate. *FASEB J*. 1999 Feb;13
- 28 Jacob S, et al. Enhancement of Glucose Disposal in Patients with Type 2 Diabetes by Alpha-lipoic Acid. *Arzneimittelforschung*. Aug1995;45(8):872-74.
- 29 Evans JL, Goldfine ID. Alpha-lipoic acid: a multifunctional antioxidant that improves insulin sensitivity in patients with type 2 diabetes. *Diabetes Technol Ther*. Sep2000;2(3):401-13.
- 30 Estrada DE, Ewart HS, Tsakiridis T, et al. Stimulation of glucose uptake by the natural coenzyme alpha-lipoic acid/thioctic acid: participation of elements of the insulin signaling pathway. *Diabetes*. 1996;45:1798-1804.

31. Ziegler D, Schatz H, Conrad F, Gries FA. Effects of treatment with the antioxidant alpha-lipoic acid on cardiac autonomic neuropathy in NIDDM patients: a 4-month randomized controlled multicenter trial (DEKAN study). *Diabetes Care* 1997;20:369-73.
32. Kunt T, et. Al.: Alpha-lipoic acid reduces expression of vascular cell adhesion molecule-1 and endothelial adhesion of human monocytes after stimulation with advanced glycation end products. *Clin Sci (Lond)*. 1999 Jan; 96(1):75-82
33. Maquart FX, et al. Stimulation of Collagen Synthesis in Fibroblast Cultures by a Triterpene Extracted from *Centella asiatica*. *Connect Tissue Res*. 1990;24(2):107-20.
34. Tenni R, et al. Effect of the Triterpenoid Fraction of *Centella asiatica* on Macromolecules of the Connective Matrix in Human Skin Fibroblast Cultures. *Ital J Biochem*. 1988;37(2):69-77.
35. Shukla A, Rasik AM, Jain GK, Shankar R, et al. In Vitro and In Vivo Wound Healing Activity of Asiaticoside Isolated from *Centella asiatica*. *J Ethnopharmacol*. Apr1999;65(1):1-11.
36. Barracchini A, Franceschini N, Filippello M, et al. Leukocyanidines and Collagenases: In Vitro Enzyme Inhibition Activity. *Clin Ter*. Jul1999;150(4):275-8.
37. Cesarone MR, Laurora G, De Sanctis MT, et al. The Microcirculatory Activity of *Centella asiatica* in Venous Insufficiency. A Double-blind Study. *Minerva Cardioangiol*. Jun1994;42(6):299-304.
38. Cesarone MR, Laurora G, De Sanctis MT, et al. Activity of *Centella asiatica* in Venous Insufficiency. *Minerva Cardioangiol*. Apr1992;40(4):137-43.
39. Belcaro GV, Grimaldi R, Guidi G. Improvement of Capillary Permeability in Patients with Venous Hypertension after Treatment with TTFCA. *Angiology*. Jul1990;41(7):533-40.
40. Belcaro GV, Rulo A, Grimaldi R. Capillary filtration and ankle edema in patients with venous hypertension treated with TTFCA. *Angiology*. 1990;41:12-18.
41. Zafirov D, Bredy-Dobrova G, Litchev V, et al. Antiexudative and Capillaritonic Effects of Procyanidines Isolated from Grape Seeds (*V. Vinifera*). *Acta Physiol Pharmacol Bulg*. 1990;16(3):50-4.
42. Blazso G, Gabor M. Oedema-inhibiting Effect of Procyanidin. *Acta Physiol Acad Sci Hung*. 1980;56(2):235-40.
43. Zafirov D, et al. Antiexudative and Capillaritonic Effects of Procyanidines Isolated from Grape Seeds (*V. vinifera*). *Acta Physiol Pharmacol Bulg*. 1990;16(3):50-54.
44. Thebaut JF, Thebaut P, Vin F. Study of Endotelon in Functional Manifestations of Peripheral Venous Insufficiency. *Gaz Med France*. 1985;92:96-100.
45. Robert L, et al. The Effect of Procyanidolic Oligomers on Vascular Permeability. A Study Using Quantitative Morphology. *Pathol Biol. (Paris)*. 1990;38(6):608-16.
46. Vanscheidt W, Jost V, Wolna P, et al. Efficacy and safety of a Butcher's broom preparation (*Ruscus aculeatus* L. extract) compared to placebo in patients suffering from chronic venous insufficiency. *Arzneimittelforschung*. 2002;52(4):243-250.
47. Capelli R, Nicora M, Di Perri T. Use of extract of *Ruscus aculeatus* in venous disease in the lower limbs. *Drugs Exp Clin Res* 1988;14:277-83.
48. Lucker P, Jost V, Wolna P, et al. Efficacy and safety of ruscus extract compared to placebo in patients suffering from chronic venous insufficiency [abstract]. *Phytomedicine*. 2000;7(suppl 2):P-155.
49. Rudofsky G, Diehm C, Gruss JD, et al. Chronic venous insufficiency. Treatment with *Ruscus* extract and trimethylhesperidin chalcone [in German; English abstract]. *MMW Munch Med Wochenschr*. 1990;132:205-210.
50. Weindorf N, Schultz-Ehrenburg U. Controlled study of increasing venous tone in primary varicose veins by oral administration of *Ruscus aculeatus* and trimethylhesperidin chalcone [in German; English abstract]. *Z Hautkr*. 1987;62:28-30,35-38.
51. Cluzan RV, Alliot F, Ghabboun S, et al. Treatment of secondary lymphedema of the upper limb with CYCLO 3 FORT. *Lymphology*. 1996;29:29-35.
52. Seltzer AP. Minimizing post-operative edema and ecchymoses by the use of an oral enzyme preparation (bromelain): a controlled study of 53 rhinoplasty cases. *Eye Ear Nose Throat Mon*. 1962;41:813-817.
53. Gylling U, Rintala A, Taipale S, et al. The effect of a proteolytic enzyme combinate (bromelain) on the postoperative oedema by oral application. A clinical and experimental study. *Acta Chir Scand*. 1966;131:193-196.